

Experimental Summary on Hadronic Decays: A TAU98 Review *

Brian K. Heltsley

^aWilson Synchrotron Laboratory, Cornell University
Ithaca, NY 14853 USA
bkh@mail.lns.cornell.edu

Selected results on hadronic decays of the tau lepton from the TAU98 Workshop are reviewed. A comprehensive picture emerges for strange particle branching fractions, and exploration of resonant substructure of both strange and non-strange decays is seen to have matured substantially.

1. Introduction

Unique among the leptons, the tau is massive enough to decay semi-hadronically. Many hadronic branching fractions are predicted with precision in the Standard Model, motivating correspondingly accurate measurements as probes of new physics. Tau decays also furnish a remarkably potent production source for mesons, the properties of which are then available for study.

Although the internal inconsistencies and disagreements between measured and predicted branching fractions have diminished in the past several years, the process of rate measurements for every conceivable mode and unrelenting improvement of precision continues. The larger branching fractions are measured with relative errors below one percent, signals for rare decays are seen at the 10^{-4} level, and forbidden decay rate upper limits approach 10^{-6} . Anticipating the completion of the LEP I and CLEO II programs, experimenters are exploiting the large datasets and pursuing improved understanding of the detectors and better analysis techniques to further study hadronic tau decays.

In earlier Workshops the experimental emphasis initially focussed on putting π^0 detection on par with that of charged particles for modes with one or multiple π^0 's; strange particle identification efforts moved from detached vertex finding for K_S , to dE/dx or RICH calibration for K^- , to calorimetry for K_L . These techniques have

been refined while datasets have grown, allowing more challenging modes involving *multiple* kaons and/or *multiple* π^0 's to be measured.

In order to make quantitative conclusions from the large number of new and old measurements, the common practice of forming weighted averages of all branching fractions is followed here. For each mode, the "New World Average" (NWA) weights each of n measurements by its inverse-square total error, which is the quadrature sum of statistical and systematic contributions. The NWA error is computed as the inverse of the quadrature sum of the inverse total errors, scaled up by the Particle Data Group [1] scale factor $\max(1, S)$, where $S = \sqrt{\chi^2/(n-1)}$. For this review, I have omitted any entries that, by virtue of their errors relative to others, contribute less than 5% of the weight in the NWA. PDG98 [1] contains the complete listings. These NWA's only are only approximate for a number of reasons: many newer measurements are preliminary; some systematic errors may be correlated and therefore should not be combined in quadrature; biased weighting can occur for statistics-limited measurements. In addition, measurements of the same quantity sometimes make different assumptions for cross-feed branching ratios or resonant content, and hence should be converted to a common basis before being averaged. Finally, a set of measurements can be internally inconsistent (have a scale factor S significantly larger than unity), in which case the NWA does not have a clear interpretation.

The scope of this summary does not allow for

*Invited talk at *The Fifth Workshop on Tau Lepton Physics*, Santander, Spain, 14-18 September 1998.

a complete or comprehensive compilation of every new result since TAU96; in particular, measurements inclusive of neutrals have been largely neglected. The reader is referred to the other presentations at this conference for up-to-date information on V/A spectral functions and the determination of α_s in tau decay [2,3], 3π sub-structure [4], and the impact of tau decays on hadronic contributions to $(g-2)_\mu$ and α_{QED} [5].

2. PDG98 branching fractions & fit

There have been several hadronic branching fraction results published since TAU96, so consulting PDG98 [1] is quite worthwhile. A systematic compilation with averages by specific decay appears along with a global fit. In the fit, measured decays are expressed in terms of a set of exclusive basis decay modes, and the sum of the basis mode branching fraction is constrained to unity. The key observation by the PDG for tau decays in this edition is that there was only one basis mode ($\mathcal{B}(\pi\nu_\tau)$) that changed from PDG96 [6] by more than one standard deviation. Here the most important feature of tau branching fractions is one of gradually improving precision, internal consistency, and stability of results. “Deficits” or “problems” in topological branching fraction vs. itemized decay mode sums, or significant internal disagreements in individual modes are mostly “old news”. The usually more precise recent measurements tend to be consistent with each other, and the troubles of the past seem to have been caused by systematic errors in the older, (now) less precise measurements.

Nevertheless it may be useful to probe the limitations of the PDG98 conclusions. The global fitting procedure does not handle explicit correlations between measured branching fractions in different modes, even though such correlations have sometimes been given along with the measurement results. Six measured decay modes totaling $(0.15 \pm 0.05)\%$ are explicitly ignored in the fit because they cannot be expressed in terms of the basis modes. A special unofficial fit with a catch-all unconstrained “dummy” decay mode results in this mode being attributed with $(0.50^{+0.40}_{-0.29})\%$ branching fraction [7]. Hence the



basis mode averages sum to less than unity, and in the official global fit, some branching fractions are pulled noticeably higher to compensate. There may be unmeasured modes with branching fraction sum at the $\sim 0.5\%$ level, or there may be problems with some of the measurements, or this may be a statistical fluctuation. However, there are other odd features to the global fit that have led the PDG to plan for a new fitter [7], one that will better address the stated goal of PDG to attain 0.1% internal consistency.

3. CVC Tests

Those tau decays that proceed through the vector current are subject to tests of the Conserved Vector Current and therefore comparison with electron-positron annihilation to the same final states. With the increase in amount and quality of tau data, in recent years tests of CVC have become limited by both the lack of comparably precise data from low energy e^+e^- colliders and the subtlety of systematic effects such as radiative corrections, separation of isoscalar components, and interference. This situation is improving somewhat with recent data from the CMD2 and SND experiments operating at the VEPP-2M collider at BINP in Novosibirsk. A compilation [8] of the measured versus predicted branching fractions for vector-current modes shows a 2.4 standard deviation excess in the overall sum of tau rates compared with CVC prediction, with the excess concentrated in the two modes with largest branching fraction: $h\pi^0$ and $3h\pi^0$ (see Figs. 2, 7). At this point it is too soon to tell whether this is significant given the experimental and theoretical subtleties involved. Further work is necessary, and improved data are expected on both tau and e^+e^- sides.



4. Non-strange Decays

OPAL [9,10] and DELPHI [11] have new branching fraction results in several non-strange modes made possible by new understanding of how π^0 's can be detected in their calorimeters. For $\mathcal{B}(h\nu_\tau)$, $\mathcal{B}(h\pi^0\nu_\tau)$, $\mathcal{B}(h2\pi^0\nu_\tau)$, $\mathcal{B}(3h^\pm\nu_\tau)$, $\mathcal{B}(5h^\pm\nu_\tau)$, and $\mathcal{B}(5h^\pm\pi^0\nu_\tau)$, as shown in Fig. 1-



Figure 1. Branching fractions for $\tau^- \rightarrow h^- \nu_\tau$.Figure 2. Branching fractions for $\tau^- \rightarrow h^- \pi^0 \nu_\tau$.

6, the new entries are compatible with and comparable in precision to previous measurements. However, the preliminary DELPHI measurement of $\mathcal{B}(3h^\pm \pi^0 \nu_\tau)$ (Fig. 7) is 2.5 standard deviations higher than the PDG98 fit value. It is also 2.6 standard deviations from the CVC prediction for $\mathcal{B}(3\pi^\pm \pi^0 \nu_\tau) = (3.90 \pm 0.22)\%$ [8] augmented by the NWA's for $\mathcal{B}(K^- \pi^+ \pi^- \pi^0 \nu_\tau) + \mathcal{B}(K^- K^+ \pi^- \pi^0 \nu_\tau)$ (Figs. 12 and 17).

There have also been new results on other high multiplicity final states. CLEO finalized their TAU96 analysis [12] on $\tau \rightarrow f_1 \pi^- \nu_\tau$ with a publication [13], reporting $\mathcal{B}(3\pi^\pm \eta \nu_\tau) = (3.4^{+0.6}_{-0.5} \pm$

Figure 3. Branching fractions for $\tau^- \rightarrow h^- 2\pi^0 \nu_\tau$.Figure 4. Branching fractions for $\tau^- \rightarrow 3h^\pm \nu_\tau$.



$0.6) \times 10^{-4}$, $\mathcal{B}(\pi^- 2\pi^0 \eta \nu_\tau) = (1.4 \pm 0.6 \pm 0.3) \times 10^{-4}$, and $\mathcal{B}(f_1 \pi^- \nu_\tau) = (5.8^{+1.4}_{-1.3} \pm 1.8) \times 10^{-4}$. A new measurement [14] of the $f_1(1285) \rightarrow \eta \pi \pi$ branching fraction as $(52.8 \pm 3.8 \pm 2.5)\%$ allows reduction of the CLEO systematic error, resulting in $\mathcal{B}(f_1 \pi^- \nu_\tau) = (5.9^{+1.4}_{-1.3} \pm 1.2) \times 10^{-4}$. In a related mode, CLEO reports $\mathcal{B}(3\pi^\pm 3\pi^0 \nu_\tau) = (2.85 \pm 0.56 \pm 0.51) \times 10^{-4}$, finding that it is saturated by the above $3\pi\eta$ modes and the (new) $\pi^- 2\pi^0 \omega$ final state with rate $\mathcal{B}(\pi^- 2\pi^0 \omega \nu_\tau) = (1.89^{+0.74}_{-0.67} \pm 0.40) \times 10^{-4}$ [15,16]. The older ALEPH measurement of $\mathcal{B}(3\pi^\pm \geq 3\pi^0 \nu_\tau) = (11 \pm 4 \pm 5) \times 10^{-4}$ [1] is larger but consistent with the new CLEO result; although it should be noted that $\mathcal{B}(3\pi^\pm \geq 4\pi^0 \nu_\tau)$ has not yet been measured directly. Measured six pion branching fractions (for $3\pi^\pm 3\pi^0$ and $5\pi^\pm \pi^0$) are found to be consistent with isospin symmetry

Figure 5. Branching fractions for $\tau^- \rightarrow 5h^\pm \nu_\tau$.Figure 6. Branching fractions for $\tau^- \rightarrow 5h^\pm \pi^0 \nu_\tau$.

once the axial-vector $(3\pi)^-\eta$ contributions have been subtracted [15,16]. Finally, new upper limits on tau decays with seven charged particles plus neutrals have been made by OPAL [17] (1.8×10^{-5} at 95%CL) and CLEO [18] (2.4×10^{-6} at 90%CL).

5. Modes involving strange particles

Much of the focus of new work on hadronic decays focused on modes with strange particles. ALEPH presented a comprehensive analysis [19] of 27 different modes, almost always measuring both K_S and K_L for modes with K^0 's.

Figure 7. Branching fractions for $\tau^- \rightarrow 3h\pi^0 \nu_\tau$.Figure 8. Branching fractions for $\tau^- \rightarrow K^- \nu_\tau$.

5.1. $\tau \rightarrow K^- \nu_\tau$

Both ALEPH [19,20] and DELPHI [21,22] have new branching ratios for $\mathcal{B}(K^- \nu_\tau)$; they are consistent with previous measurements, and ALEPH halves their previous (smallest) error, which improves the world average accordingly, as shown in Fig. 8. The ALEPH analysis uses this improved precision to test $\tau - \mu$ universality ($g_\tau/g_\mu = 0.987 \pm 0.021$) [23] by comparing the decay rate to that of muonic kaon decay, and to determine the kaon decay constant, assuming lepton universality, as $f_K = (111.5 \pm 0.9 \pm 2.3)$ MeV.

5.2. $\tau \rightarrow (K\pi)^- \nu_\tau$

ALEPH [19,20,23] presented new results for $K^-\pi^0$, $K_S\pi^-$, and $K_L\pi^-$ final states. Fig. 9 shows the $(K\pi)^-$ branching fraction, labeled by specific final state, assuming an isospin $\frac{1}{2}$ parent.




Figure 9. Branching fractions for combined $\tau^- \rightarrow (K\pi)^-\nu_\tau$, assuming an isospin $\frac{1}{2}$ parent such as $K^*(892)$ and/or $K^*(1410)$.

The CLEO result remains about two standard deviations lower than the more precise combined ALEPH rate.

ALEPH also fits the $K\pi$ spectrum for the presence of $K^*(1410)$ in addition to $K^*(892)$, and for interference between the two final states. They obtain evidence for $K^*(1410)$ at the one to three standard deviation level, depending upon which parameters are fixed or allowed to float in the fit and which decay mode spectra are included. The $K^*(1410)$ admixture strength β is determined to be ~ -0.1 , similar to that measured for $\rho'(1450)$ in $\tau \rightarrow \pi^-\pi^0\nu_\tau$ [24,25]. The net branching fraction is measured as $\mathcal{B}(K^*(1410)\nu_\tau) = (0.15^{+0.14}_{-0.10})\%$. As $K^*(1410)$ decays 93% of the time to $(K\pi\pi)^-$, this measurement yields the vector component background to the dominantly axial-vector $(K\pi\pi)^-$ final states.

5.3. $\tau \rightarrow (K\pi\pi)^-\nu_\tau$

In analogy with tau decays to non-strange $(3\pi)^-$ states, the production of $(K\pi\pi)^-$ is expected to occur through the axial-vector mesons $K_1(1270)$ and $K_1(1400)$ [26–30]. However, immediate non-resonant fragmentation to $(K^*\pi)^-$ or $(K\rho)^-$ at the pointlike weak vertex, or production through the strange vector meson $K^*(1410)$ (see previous section), are also possible.

Let K_A and K_B denote the triplet 3P_1 and singlet 1P_1 spin eigenstates, respectively, of $(\bar{u}s)$, the strange analogs of the $\bar{u}d$ states, $a_1(1260)$ and $b_1(1235)$. Production amplitudes of b_1 and K_B in tau decay are suppressed in the SM as second-class currents by

$$\sim \frac{|m_u - m_{d,s}|}{m_u + m_{d,s}}.$$

This factor is quite small for m_d , accounting for the non-observation of b_1 in tau decays. In contrast, the large strange quark mass induces substantial $SU(3)_f$ symmetry-breaking, allowing non-zero K_B production in tau decay.

The strange axial-vector mass eigenstates are $K_1(1270)$ and $K_1(1400)$. $K_1(1270)$ decays to $K\rho$, $K_0^*(1430)\pi$, $K^*\pi$, and $K\omega$; $K_1(1400)$ decays almost exclusively to $K^*\pi$. Since K_A and K_B are both predicted by $SU(3)$ to decay equally to $K^*\pi$ and $K\rho$, and since the K_1 states do not, the K_1 resonances must correspond to mixtures of K_A and K_B in comparable amounts with opposite-sign phases. One can parametrize this mixing in terms of an angle θ_K :

$$K_1(1400) = K_A \cos\theta_K - K_B \sin\theta_K$$

$$K_1(1270) = K_A \sin\theta_K + K_B \cos\theta_K$$

The previously measured properties of the K_1 mesons place some restrictions on the allowed values of θ_K [27,28]. The relative amounts of $K_1(1270)$ and $K_1(1400)$ observed in tau decay can further constrain θ_K and yield the size of $SU(3)_f$ symmetry-breaking.

ALEPH [19,20,23] presented new information on all four possible $K\pi\pi$ final states, while CLEO [31] and DELPHI [21] did so for the all-charged mode. The updated ALEPH value $\mathcal{B}(K^-2\pi^0\nu_\tau) = (0.056 \pm 0.025)\%$ is now corrected for $K^-3\pi^0$ feed-down. Fig. 10 compare the CLEO and ALEPH results for the K^0 modes, which are consistent, and Fig. 11 does the same for the $K^-\pi^+\pi^-$ mode, for which the three results differ substantially from one another. The theoretical prediction from Li [32] of 0.18% is closer to the ALEPH and CLEO result than Finkemeier and Mirkes [33] prediction of 0.77%, which is much higher even than the DELPHI measurement.




Figure 10. $\tau^- \rightarrow \bar{K}^0 \pi^- \pi^0 \nu_\tau$ branching fractions.




Figure 11. $\tau^- \rightarrow K^- \pi^+ \pi^- \nu_\tau$ branching fractions.

CLEO [31] and ALEPH [19,23] have both made quantitative analyses of the resonance structure of $(K\pi\pi)^-$ decays. CLEO has $\sim 8K$ events in $K^-\pi^+\pi^-$ with 38% background and is systematics-limited, while ALEPH has ~ 300 events split between the $K^-\pi^+\pi^-$ and $\bar{K}^0\pi^-\pi^0$ modes, each with $\sim 50\%$ background. CLEO fits the sub-mass spectra for $K_1(1270)$, $K_1(1400)$ amplitudes, strong phases in their $K^-\rho$ and $K^{*0}\pi^-$ decays, and the mixing angle θ_K . Accounting for the interference is found to be important, as the two phases are about 5 and 1.7 standard deviations from zero: the predicted mass spectra match the data well only if these phases are included. The $K_1(1270)$ is found to dominate, and CLEO quotes the ratio of the two K_1 states as $K_1(1270)/K_1(1400) = 4.1 \pm 0.5^{+4.8}_{-2.1}$. The final state $K\rho$ is found to be $(45 \pm 12)\%$ of $K^-\pi^+\pi^-$.

Finally, the mixing angle is determined within a two-fold ambiguity to be $(67^\circ \text{ or } 46^\circ) \pm 10^\circ$, close to the values obtained from K_1 widths and branching fractions [27] of $(57^\circ \text{ or } 33^\circ)$.

In contrast, ALEPH assumes the $K\pi\pi$ state appears as an *incoherent* superposition of $K\rho$ and $K^*\pi$, finding $K\rho$ to be $(35 \pm 11)\%$ of $K^-\pi^+\pi^-$ and $(66 \pm 12)\%$ of $\bar{K}^0\pi^-\pi^0$ (consistent with isospin symmetry), the fraction of $K_1(1270)$ in the final states to be $(41 \pm 19 \pm 15)\%$, and the branching fraction $\mathcal{B}(K_1(1270)\nu_\tau) = (0.45 \pm 0.11)\%$. So, while ALEPH and CLEO use different assumptions and analyze their data differently, qualitatively, they agree that $K\rho$ is a significant presence in $K\pi\pi$ decays, and therefore that $K_1(1270)$ plays a major role as a source of these final states.

Another final state may proceed through the K_1 resonance. CLEO [34] has found $\mathcal{B}(K^{*-}\eta\nu_\tau) = (2.9 \pm 0.8 \pm 0.4) \times 10^{-4}$, finding that the K^* is dominant in $(K\pi)^-\eta$ decays. Li [32] predicts that this final state proceeds through the K_1 axial-vector current, and, using an effective chiral theory in the limit of chiral symmetry, predicts a branching fraction of (1.01×10^{-4}) . In contrast, Pich [35] predicts a rate for $(K\pi)^-\eta$ of $\sim 10^{-5}$ using chiral perturbation theory.

5.4. $\tau \rightarrow (K\pi\pi\pi)^-\nu_\tau$

ALEPH [19,23] and CLEO [31] both look for $(K3\pi)^-$ final states, do not obtain definitive signals for any of them, but determine sub-0.1% branching fractions. Fig. 12 shows the measurements for $K^-\pi^+\pi^-\pi^0$, which are consistent with each other. ALEPH also measures $\mathcal{B}(\bar{K}^0\pi^-2\pi^0(excl. K_S)\nu_\tau) = (0.058 \pm 0.036)\%$, $\mathcal{B}(K^-3\pi^0(excl. \eta)\nu_\tau) = (0.037 \pm 0.024)\%$, and $\mathcal{B}(K^03h^\pm(excl. K_S)\nu_\tau) = (0.023 \pm 0.020)\%$.

5.5. Total Strange Spectral Function

ALEPH [3] has combined the spectral information from $K\pi$ and $K\pi\pi$ decays with small additions from Monte Carlo for $K \geq 3\pi$ modes to determine the total strange spectral function. Using this and the strange-quark mass dependence of the QCD expansion for the hadronic decay width of the tau, ALEPH obtains $m_S(1 \text{ GeV}^2) = (217^{+45}_{-57}) \text{ MeV}$.




Figure 12. $\tau^- \rightarrow K^-\pi^+\pi^-\pi^0\nu_\tau$ branching fractions.




Figure 13. Branching fractions for $\tau^- \rightarrow K^0 K^-\nu_\tau$.

5.6. $\tau \rightarrow (KK)^-\nu_\tau, (KK\pi)^-\nu_\tau, (KK\pi\pi)^-\nu_\tau$

The production of $(KK)^-$ or $(KK\pi)^-$ states in tau decay is suppressed by phase space, but could occur via direct fragmentation or through the high-mass tails of the ρ^- , a_1^- , or ρ'^- : $\rho^- \rightarrow (KK)^-$, $\rho'^- \rightarrow (K^*K)^-$, $a_1^- \rightarrow (\rho\pi)^- \rightarrow (KK\pi)^-$, or $a_1^- \rightarrow (K^*K)^-$. ALEPH and CLEO have new results in many of these modes, including $K^-K^+\pi^-\pi^0$, as shown in Figs. 13-17. There are no major inconsistencies in the measured branching fractions, which are typically $\sim 0.1\%$. The observation by CLEO of K^*K content in $3h^\pm$ events is quantitatively consistent with the measured branching ratio: that analysis [4,36] determines $\mathcal{B}(\tau \rightarrow K^*K\nu_\tau)/\mathcal{B}(\tau \rightarrow (3h)^-\nu_\tau) = (3.3 \pm 0.5 \pm 0.1)\%$, which results in the prediction for the $K^-K^+\pi^-$ branching fraction shown in Fig. 14.

The ALEPH spectrum for $K_S K^-$ does not match the model expectation for $\rho \rightarrow K\bar{K}$, in




Figure 14. $\tau^- \rightarrow K^-K^+\pi^-\nu_\tau$ branching fractions.




Figure 15. $\tau^- \rightarrow K_S^0 \bar{K}_S^0 \pi^-\nu_\tau$ branching fractions.

agreement with the conclusion of CLEO [37]. The ALEPH analysis also favors K^*K parentage over $\rho\pi$ in $KK\pi$ states, and quotes an axial-vector current fraction of $(94_{-8}^{+6})\%$ based on CVC and the absence of $KK\pi$ in e^+e^- data.

Some comments on the $K^0 \bar{K}^0 \pi^-$ final state [1] are in order. ALEPH sees 68 $K_S K_L \pi^-$ events and 6 in $K_S K_S \pi^-$, compared to CLEO's 52 events [37] in the latter mode only. If the K^0 and $\bar{K}^0$ decay independently, the $K_S K_L \pi^-$ mode should have double the rate of $K_S K_S \pi^-$ or $K_L K_L \pi^-$. However, Bose-Einstein correlations between the neutral kaons will violate this assumption. The ALEPH measurement [19] of $\mathcal{B}(K_S K_L \pi^- \nu_\tau) = (0.101 \pm 0.023 \pm 0.013)\%$ is two standard deviations higher than double the NWA $\mathcal{B}(K_S K_S \pi^- \nu_\tau)$, suggestive that such an effect may be in force, but errors are currently too high for any definitive conclusion.




Figure 16. $\tau^- \rightarrow K^0 K^- \pi^0 \nu_\tau$ branching fractions.




Figure 17. $\tau^- \rightarrow K^- K^+ \pi^- \pi^0 \nu_\tau$ branching fractions.

6. Conclusions

Some perspective on recent experimental accomplishments can be gained by answering some of the questions I posed in the TAU94 proceedings [30]: *Will the global branching fractions of ALEPH continue to dominate most modes, or can we expect comparably precise results from other experiments?* ALEPH still sets the experimental standard for comprehensive treatment of hadronic tau decays, due to a combination of the clean tau-pair topology at LEP energies, a superbly designed and maintained detector, hard work, and aggressive analysis. This is particularly true of decays involving a K_L , or a charged kaon and π^0 's, or modes with multiple π^0 's. CLEO competes best on modes with π^0 's and/or with branching fractions below 1% where its large dataset can provide enough events. In a very

welcome development, OPAL and DELPHI have recently achieved better understandings of their detector systematics and are pressing or even surpassing ALEPH and CLEO on selected modes. It is a trend we hope can continue. However, we all await ALEPH's final full-data-sample global analysis with trepidation!

Can we resolve branching fraction discrepancies in the $h\pi^0$ and $3h$ modes simply by improving measurements, making older, disagreeable values irrelevant? Apparently so.

Can we develop procedures which allow a consistent accounting of modes with K_L decays? PDG96 [6] adopted basis modes which excluded unseen K_L , and measurements of modes with K_L , dominated by ALEPH [19], have made the feed-across corrections feasible.

What are the masses and widths of the ρ and ρ' in tau decay? ALEPH results have been published [24], but the preliminary results from CLEO [25] have not.

Is all $(3\pi)^-$ attributable to a_1^- decays? A poorly-phrased question; better would have been “are there intermediate states in $(3\pi)^-$ that do not go through $a_1 \rightarrow \rho\pi$?”. The CLEO analysis [4,36] shows that less than 70% of $\pi^-\pi^0\pi^0$ final states come from $\rho\pi$ intermediate states, and in particular that isoscalars ($\sigma\pi$, $f_0(1186)\pi$) amount to $\sim 20\%$ of the branching fraction. The CLEO and DELPHI [38] analyses also are suggestive of the presence of an a_1' at high 3π mass.

Is there evidence for scalar or non-resonant production of $(K\pi)^-$ final states? No, but statistics are scarce for this study. I should have asked “is there evidence for $K^*(1410)$ in $K\pi$ decays”, to which the answer is, yes, from ALEPH [19,23].

Are $(K\pi\pi)^-$ decays consistent with a mixture of $K_1(1270)$ and $K_1(1400)$ production? Yes, and both CLEO [31] and ALEPH [19] agree that $K_1(1270)$ amounts to a substantial fraction of the $(K\pi\pi)^-$ final state. *What is the mixing angle and strength of $SU(3)_f$ symmetry breaking in such decays?* CLEO has a preliminary analysis [31], which accounts for interference between the different intermediate states, in which the mixing angle is measured up to a two-fold ambiguity. *Do $(K\pi\pi)^-$ final states have $K^*(1410)$ or non-resonant content?* ALEPH [19,23] has

measured $\mathcal{B}(K^*(1410)\nu_\tau) = (0.15^{+0.14}_{-0.10})\%$, and so the branching fraction into $(K\pi\pi)^-$ can be computed. Statistics are too low to answer for non-resonant production.

Can all $(4\pi)^-$ final states be described as $\omega\pi^-$ and $(a_1\pi)^-$? If so, what resonances produce those states? A complete spectral analysis of $\tau^- \rightarrow (4\pi)^-\nu_\tau$ has yet to be done.

What is the resonant substructure in $KK\pi$, 5π , 6π final states? The $KK\pi$ final state seems to arise through the axial-vector current $a_1^- \rightarrow K^*K$ [19,23]. Five and six pion final states are quite complex, but receive important contributions from $f_1\pi$ and $\pi^-2\pi^0\omega$ [13,15,16].

What is the rate for 7π production in tau decay? For seven charged pions with or without π^0 's, less than 2.4 parts per million [15,18]. No direct measurement of $3\pi^\pm 4\pi^0$ has been performed and only upper limits exist for $5\pi^\pm 2\pi^0$.

Is there any evidence for second-class currents in tau decay at expected or unexpected levels? Sadly, nothing unexpected, and the expected levels have yet to be approached experimentally.

Given the current status of measurements, where are the most promising places to look for deviations from the Standard Model in hadronic tau decays? Recent effort has been focussed on rare, suppressed, or forbidden decays [15], and on CP-violation [39], but any Standard Model prediction is fair game.

Looking ahead, as the LEP and CLEO II programs draw to a close, finalized, full-statistics data-sets should become available next year for analysis. Improved low-energy data for CVC tests should continue to emerge from the experiments at VEPP-2M. Considerably more hadronic tau physics awaits further mining of all these samples, both with high-statistics systematics-limited studies which need more time and data to mature, and for low-statistics searches which can benefit the most from additional luminosity. Experimenters should not be dissuaded from this pursuit by the somewhat disquieting over-consistency of many results; surprise often lurks just beyond the horizon. We should all look forward to a peek at something new at TAU2000.

7. Acknowledgements

I would like to thank other speakers for making their results available to me for the writing of this summary talk. In particular, I benefitted greatly from communications with S. Chen, M. Davier, S. Eidelman, J.M. Lopez-Garcia, and R. Sobie. My fellow CLEOns continue to indulge my endless questions, especially J. Smith, I. Kravchenko, D. Besson, and A. Weinstein. The Organizers and Secretariat are to be commended for hosting a productive and invigorating Workshop. The preparation of this manuscript was supported by the National Science Foundation.

REFERENCES

1. C. Caso *et al.* (Particle Data Group) Eur. Phys. J. C **3** (1998) 286.
2. S. Menke, presentation at this Workshop.
3. A. Hoecker, presentation at this Workshop.
4. M. Schmidtler, presentation at this Workshop.
5. M. Davier, presentation at this Workshop.
6. R.M. Barnett *et al.* (Particle Data Group) Phys. Rev. D. **54** (1996) 256.
7. K. Moenig, private communication.
8. S. Eidelman, presentation at this Workshop.
9. K. Ackerstaff *et al.* (OPAL Collaboration) Eur. Phys. J. C **4** (1998) 193.
10. K. Ackerstaff *et al.* (OPAL Collaboration) CERN-EP/98-090 (June 1998) (Submitted to Eur. Phys. J.).
11. J.M. Lopez, presentation at this Workshop.
12. V. Shelkov, in *Proceedings of the Fourth Workshop on Tau Lepton Physics*, September 1996, edited by J.G. Smith and W. Toki, Nucl. Phys. B (Proc. Suppl.) **55C** (1997) 195.
13. T. Bergfeld *et al.* (CLEO Collaboration), Phys. Rev. Lett. **79** (1997) 2406.
14. D. Barberis *et al.* (WA102 Collaboration), hep-ex/9810003 (September 1998).
15. K.K. Gan, presentation at this Workshop.
16. S. Anderson *et al.* (CLEO Collaboration), Phys. Rev. Lett. **79** (1997) 3814.
17. K. Ackerstaff *et al.* (OPAL Collaboration), Phys. Lett. **B404** (1997) 213.
18. K. Edwards *et al.* (CLEO Collaboration),

- Phys. Rev. D **56** (1997) 5297.
19. S. Chen, presentation at this Workshop.
 20. R. Barate *et al.* (ALEPH Collaboration), ALEPH 98-083 (September 1998).
 21. A. Andreazza, presentation at this Workshop.
 22. DELPHI Collaboration, DELPHI 98-145 & ICHEP98 #247, (June 1998).
 23. R. Barate *et al.* (ALEPH Collaboration), ALEPH 98-084 (September 1998).
 24. R. Barate, *et al.* (ALEPH Collaboration), Z. Phys. **C76** (1997) 15.
 25. J. Urheim, in *Proceedings of the Fourth Workshop on Tau Lepton Physics*, September 1996, edited by J.G. Smith and W. Toki, Nucl. Phys. B (Proc. Suppl.) **55C** (1997) 359.
 26. M.T. Ronan, in *Proceedings of the Second Workshop on Tau Lepton Physics*, September 1992, edited by K.K. Gan, World Scientific (1993), p. 225.
 27. M. Suzuki, Phys. Rev. D**47** (1993) 1252.
 28. M. Suzuki, Phys. Rev. D**49** (1994) 2634.
 29. H.J. Lipkin, Phys. Lett. **B303** (1993) 119.
 30. B. Heltsley, in *Proceedings of the Third Workshop on Tau Lepton Physics*, September 1994, edited by L. Rolandi, Nucl. Phys. B (Proc. Suppl.) **40** (1995) 413.
 31. I. Kravchenko, presentation at this Workshop.
 32. B.A. Li, Phys. Rev. D **55** (1997) 1436; Phys. Rev. D **57** (1998) 1790.
 33. M. Finkemeier and E. Mirkes, Z. Phys. **C69** (1996) 243.
 34. M. Bishai *et al.* (CLEO Collaboration), CLNS 98/1574 (hep-ex/9809012) (1998).
 35. A. Pich, Phys. Lett. **B196** (1987) 561.
 36. T.E. Coan *et al.* (CLEO Collaboration), ICHEP98 #976 (July 1998).
 37. T. Coan *et al.* (CLEO Collaboration), Phys. Rev. D **53** (1996) 6037.
 38. P. Abreu *et al.* (DELPHI Collaboration), Phys. Lett. **B426** (1998) 411.
 39. R. Kass, presentation at this Workshop.